

UNIVERSITI TEKNOLOGI MARA

**HPLC METHOD DEVELOPMENT &
STRUCTURAL ELUCIDATION
OF ANTI-PROLIFERATIVE
QUASSINOIDS FROM
Quassia borneensis NOOT.
(SIMAROUBACEAE)**

JULENAH BINTI AG NUDDIN

Thesis submitted in fulfilment
of the requirements for the degree of
Doctor of Philosophy

Faculty of Applied Sciences

May 2016

CONFIRMATION BY PANEL OF EXAMINERS

I certify that a panel of examiners has met on 16th of June 2015 to conduct the final examination of Julenah Ag Nuddin on her Doctor of Philosophy thesis entitled "HPLC Method Development and Structural Elucidation of Anti-proliferative Quassinoids from *Quassia borneensis* Noot. (Simaroubaceae)" in accordance with Universiti Teknologi MARA Act 1976 (Akta 173). The Panel of Examiners recommends that the student be awarded the relevant degree. The Panel of Examiners was as follows:

Nor Hadiani Ismail, PhD
Professor
Faculty of Applied Sciences
Universiti Teknologi MARA
(Chairman)

Rohaya Ahmad, PhD
Professor
Faculty of Applied Sciences
Universiti Teknologi MARA
(Internal Examiner)

Amani Awaad, PhD
Professor
Faculty of Pharmacy
University of Prince Sattam bin Abdul Aziz
(External Examiner)

Hasnah Osman, PhD
Professor
School of Chemical Sciences
Universiti Sains Malaysia
(External Examiner)


:

DR MOHAMMAD NAWAWI
DATO' HAJI SEROJI
Dean
Institute of Graduate Studies
Universiti Teknologi MARA
Date : 3rd May, 2016

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name	:	Julenah binti Ag Nuddin
Student I.D. No.	:	200988294
Programme	:	Doctor of Philosophy (Chemistry) (AS990)
Faculty	:	Applied Sciences
Thesis Title:		HPLC Method Development and Structural Elucidation of Anti-proliferative Quassinoids from <i>Quassia borneensis</i> Noot. (Simaroubaceae)
Signature of Student	:	
Date	:	May 2016

ABSTRACT

Bitter quassinoids have been reported to possess numerous biological activities. They are degraded triterpenoids classified according to their basic skeletons; C₁₈, C₁₉, C₂₀, C₂₂, C₂₅. Currently, more than 200 quassinoids have been isolated from genus *Quassia*, *Brucea*, *Soulamea*, *Eurycoma*, *Picrolemania* and others. A study on *Quassia borneensis* Noot. (Simaroubaceae) (Qb) for its anti-proliferative active quassinoids using a developed and validated high performance liquid chromatography (HPLC) isolation method is appropriate for data gathering. In addition, a total synthesis exploration of canthin-6-one would provide information regarding its mechanism and influential factors for viability study. The study was accomplished by extracting powdered bark and root Qb by cold maceration and soxhlet in methanol for yield comparison. The methanol extracts were partitioned to acquire n-hexane, chloroform and aqueous fractions. They were tested for their anti-proliferative activity based on MTS assay using HL-60 cell line. Later, chloroform and aqueous fractions were subjected to isolation and purification with HPLC. Then, pure isolates were structurally elucidated with liquid chromatography-mass spectroscopy (LC-MS) and nuclear magnetic resonance (NMR). In addition, the mass profile of active fractions was measured using liquid chromatography-mass spectrometer triple quadrupole time of flight (LC-MSQTOF). Lastly, pure quassinoid compounds were subjected to MTT assay. In a biomimetic total synthesis attempt, canthin-6-one of Simaroubaceae was pursued through carboxyl-mediated approach following the method of Czerwinski et al. (2003). The positively identified sample was collected at Tawai Forest Reserve, Telupid, Sabah, Malaysia. Qb was successfully extracted and partitioned. Later, the anti-proliferative activity of hexane (Q₁-Q₄), chloroform (Q₅-Q₈) and aqueous (Q₉-Q₁₂) fractions were found active except Q₂ while Q₈ was found to be most active at 0.2 µg/mL. HPLC isolation method was developed and validated in a chromatographic system consists of Zorbax SB C-18 250 mm x 4.6 mm, 5 µm stationary phase with H₂O:ACN as its mobile phase flowing at 0.6 mL/min in step gradient elution mode of 0-100-0 % ACN with 10 % increment in an interval of five-minute for 50 minutes and 50% decrement twice in 10 minutes as post equilibration. The acquired values of selectivity factor (α), retention factor (k'), number of theoretical plates (N) and resolution (R) are at optimum indicating good separation and column performance. A peak at retention time of 21.7 minute was chosen which gives standard deviation of $\sigma=0.04$ at N=5, indicating chromatographic system usability. Thus, Q₈ was fractionated due to its best HPLC profile and anti-proliferative activity. Five quassinoids and one canthin-6-one were successfully structurally elucidated and identified as glaucarubolone (**10**), chaparrinone (**8**), holacanthone (**9**), glaucarubinone (**11**), ailanthinone (**102**) and canthin-6-one (**7**) in reference to past reports. The mass profile of chloroform fractions implied that the root part contains more quassinoids while soxhlet procedure would yield more extract. Glaucarubolone, chaparrinone, holacanthone and ailanthinone were found to be significantly active at IC₅₀ of 6.8, 7.2, 5.6 and 2.8 µM respectively. As new data, ailanthinone was found to be comparably active to doxorubicin which was tested to be active at 0.9 µM. The synthesis work of canthin-6-one yielded 64.51%, 67.05%, 48.86%, 33.56% and 0.6% of Nb, Nbb, Nbc, Nbd and Nbe respectively. A schematic mechanism of PSR in carboxyl-mediated synthesis of canthin-6-one and its influential factors were discussed. It is suggested that further isolation should be performed to Qb as more quassinoids might be found.

TABLE OF CONTENTS

	Page
CONFIRMATION BY PANEL OF EXAMINERS	ii
AUTHOR'S DECLARATION	iii
ABSTRACT	iv
ACKNOWLEDGEMENT	v
TABLE OF CONTENTS	vi
LIST OF TABLES	viii
LIST OF FIGURES	xi
LIST OF PLATES	xiii
LIST OF SCHEMES	xiv
LIST OF SYMBOLS	xvi
LIST OF ABBREVIATION	xviii
LIST OF CHEMICAL COMPOUNDS	xxi
CHAPTER ONE: INTRODUCTION	1
1.1 Traditional Medicinal Plant Family of Simaroubaceae	2
1.1.1 <i>Quassia borneensis</i> Noot.	3
1.2 Quassinoids and Canthin-6-ones of Simaroubaceae	4
1.3 High Performance Liquid Chromatography in Isolation of Quassinoids	7
1.4 Total Synthesis of Canthin-6-one	9
1.5 Problem Statement & Significance of Study	10
1.6 Objectives of Study	11
CHAPTER TWO: REVIEW OF LITERATURE	13
2.1 The Medicinal Plant Family of Simaroubaceae	13
2.1.1 <i>Quassia</i> L.	17
2.2 The Chemistry of Quassinoids	20
2.3 The Chemistry of Canthin-6-ones	27
2.3.1 Pictet-Spengler Condensation Reaction	29